

Fast convergent method for the m -point problem in Banach space

D.O. Sytnyk* V.B. Vasylyk[†]

Abstract

The m -point nonlocal problem for the first order differential equation with an operator coefficient in a Banach space X is considered. An exponentially convergent algorithm is proposed and justified provided that the operator coefficient A is strongly positive and some existence and uniqueness conditions are fulfilled. This algorithm is based on representations of operator functions by a Dunford-Cauchy integral along a hyperbola enveloping the spectrum of A and on the proper quadratures involving short sums of resolvents. The efficiency of the proposed algorithms is demonstrated by numerical examples.

Keywords nonlocal problem, differential equation with an operator coefficient in Banach space, operator exponential, exponentially convergent algorithms

AMS Subject Classification 65J10, 65M70, 35K90, 35L90

1 Introduction

The m -point initial (nonlocal) problem for a differential equation with the nonlocal condition $u(t_0) + g(t_1; \dots; t_p; u) = u_0$ and a given function g on a given point set $P = \{0 = t_0 < t_1 < \dots < t_p\}$, is one of the important topics in the study of differential equations. Interest in such problems originates mainly from some physical problems with a control of the solution at P . For

*Institute of Mathematics of NAS of Ukraine, 3 Tereshchenkivs'ka Str., Kyiv-4, 01601, Ukraine (sytnik@imath.kiev.ua).

[†]Institute of Mathematics of NAS of Ukraine, 3 Tereshchenkivs'ka Str., Kyiv-4, 01601, Ukraine (vasylyk@imath.kiev.ua).

example, when the function $g(t_1; \dots; t_p; u)$ is linear, we will have the periodic problem $u(t_0) = u(t_1)$. Problems with nonlocal conditions arise in the theory of physics of plasma [1], nuclear physics [2], mathematical chemistry [3], waveguides [4] etc.

Differential equations with operator coefficients in some Hilbert or Banach space can be considered as meta-models for systems of partial or ordinary differential equations and are suitable for investigations using tools of the functional analysis (see e.g. [5, 6]). Nonlocal problems can also be considered within this framework [7, 8].

In this paper we consider the following nonlocal m -point problem:

$$\begin{aligned} u'_t + Au &= f(t), \quad t \in [0, T] \\ u(0) + \sum_{k=1}^m \alpha_k u(t_k) &= u_0, \quad 0 < t_1 < t_2 < \dots < t_m \leq T, \end{aligned} \tag{1}$$

where $\alpha_k \in \mathbb{R}$, $k = \overline{1, m}$, $f(t)$ is a given vector-valued function with values in a Banach space X , $u_0 \in X$. The operator A with the domain $D(A)$ in a Banach space X is assumed to be a densely defined strongly positive (sectorial) operator, i.e. its spectrum $\Sigma(A)$ lies in a sector of the right half-plane with the vertex at the origin and the resolvent decays inversely proportional to $|z|$ at the infinity (see estimate (3) below).

Discretization methods for differential equations in Banach and Hilbert spaces were intensively studied in the last decade (see e.g. [9, 10, 11, 12, 13, 14, 15, 16, 17, 18] and the references therein). Methods from [10, 11, 12, 13, 14, 16, 17, 18] possess an exponential convergence rate, i.e. the error estimate in an appropriate norm is of the type $\mathcal{O}(e^{-N^\alpha})$, $\alpha > 0$ with respect to a discretization parameter $N \rightarrow \infty$. For a given tolerance ε such discretizations provide optimal or nearly optimal computational complexity [10]. One of the possible ways to obtain exponentially convergent approximations to abstract differential equations is based on a representation of the solution through the Dunford -Cauchy integral along a parametrized path enveloping the spectrum of the operator coefficient. Choosing a proper quadrature for this integral we obtain a short sum of resolvents. Since the treatment of such resolvents is usually the most time consuming part of any approximation this leads to a low-cost naturally parallelizable algorithms. Parameters of the algorithms from [12, 14, 16] were optimized in [19, 20] to improve the convergence rate.

The aim of this paper is to construct an exponentially convergent approximation to the problem for a differential equation with m -point nonlocal condition in abstract setting (1). The paper is organized as follows. In Section 2 we discuss the existence and uniqueness of the solution as well as its representation through input data. An algorithm for the homogeneous

problem (1) is proposed in section 3. The main result of this section is theorem 2 about the exponential convergence rate of the proposed discretization. The next section 4 is devoted to the exponentially convergent discretization of the inhomogeneous problem. In section 5 we represent some numerical examples which confirm theoretical results from the previous sections.

2 Existence and representation of the solution

Let the operator A in (1) be a densely defined strongly positive (sectorial) operator in a Banach space X with the domain $D(A)$, i.e. its spectrum $\Sigma(A)$ lies in the sector (spectral angle)

$$\Sigma = \left\{ z = \rho_0 + re^{i\theta} : r \in [0, \infty), |\theta| < \varphi < \frac{\pi}{2} \right\}. \quad (2)$$

Additionally outside the sector and on its boundary Γ_Σ the following estimate for the resolvent holds true

$$\|(zI - A)^{-1}\| \leq \frac{M}{1 + |z|}. \quad (3)$$

The numbers ρ_0, φ are called the spectral characteristics of A .

The hyperbola

$$\Gamma_0 = \{z(\xi) = \rho_0 \cosh \xi - ib_0 \sinh \xi : \xi \in (-\infty, \infty), b_0 = \rho_0 \tan \varphi\} \quad (4)$$

in turn is referred as a spectral hyperbola. It has a vertex at $(\rho_0, 0)$ and asymptotes which are parallel to the rays of the spectral angle Σ .

A convenient representation of operator functions is the one through the Dunford-Cauchy integral (see e.g. [5, 6]) where the integration path plays an important role. We choose the following hyperbola

$$\Gamma_I = \{z(\xi) = a_I \cosh \xi - ib_I \sinh \xi : \xi \in (-\infty, \infty)\}, \quad (5)$$

as the integration contour which envelopes the spectrum of A .

For an arbitrary vector $u_0 \in D(A^{m+1})$ the next equality holds

$$\sum_{k=1}^{m+1} \frac{A^{k-1}u_0}{z^k} + \frac{1}{z^{m+1}}(zI - A)^{-1}A^{m+1}u_0 = (zI - A)^{-1}u_0. \quad (6)$$

This formula, together with

$$A^{-(m+1)}v = \frac{1}{2\pi i} \int_{\Gamma_I} z^{-(m+1)}(zI - A)^{-1}v dz, \quad (7)$$

by setting $v = A^{m+1}u_0$, yields the following representation

$$\begin{aligned} u_0 &= A^{-(m+1)}A^{m+1}u_0 = \frac{1}{2\pi i} \int_{\Gamma_I} z^{-(m+1)}(zI - A)^{-1}A^{m+1}u_0 dz \\ &= \int_{\Gamma_I} \left[(zI - A)^{-1} - \sum_{k=1}^{m+1} \frac{A^{k-1}}{z^k} \right] u_0 dz. \end{aligned} \quad (8)$$

Various useful properties of this representation as well as the next result were discussed in [21].

Theorem 1 *Let $u_0 \in D(A^{m+\alpha})$ for some $m \in \mathbb{N}$ and $\alpha \in [0, 1]$, then the following estimate holds true:*

$$\left\| \left[(zI - A)^{-1} - \sum_{k=1}^{m+1} \frac{A^{k-1}}{z^k} \right] u_0 \right\| \leq \frac{1}{|z|^{m+1}} \frac{(1+M)K}{(1+|z|)^\alpha} \|A^{m+\alpha}u_0\|, \quad (9)$$

$\forall \alpha \in [0, 1], u_0 \in D(A^{m+\alpha}),$

where the constant K depends on α and M only.

In [7, 8] it was proven that the solution of the problem (1) exists and is unique provided that one of the following two conditions is fulfilled:

$$\sum_{i=1}^m |\alpha_i| < 1, \quad (10)$$

or

$$\sum_{i=1}^m |\alpha_i| e^{-\rho_0 t_i} < 1, \quad (11)$$

with ρ_0 from (2).

According to the Hille-Yosida-Phillips theorem [22] the strongly positive operator A generates a one parameter semigroup $T(t) = e^{-tA}$ and the solution of (1) can be represented by

$$u(t) = e^{-At}u(0) + \int_0^t e^{-A(t-\tau)} f(\tau) d\tau. \quad (12)$$

Combining the nonlocal condition from (1) and (12) we obtain

$$u(t_i) = e^{-At_i} \left[u_0 - \sum_{k=1}^n \alpha_k u(t_k) \right] + \int_0^{t_i} e^{-A(t_i-\tau)} f(\tau) d\tau, \quad (13)$$

$i = \overline{1, m}.$

Multiplying these equations by α_i and summing up over $i = \overline{1, m}$ one gets

$$\begin{aligned} \sum_{i=1}^m \alpha_i u(t_i) &= \sum_{i=1}^m \alpha_i e^{-At_i} u_0 - \sum_{i=1}^m \alpha_i e^{-At_i} \sum_{k=1}^m \alpha_k u(t_k) \\ &\quad + \sum_{i=1}^m \alpha_i \int_0^{t_i} e^{-A(t_i-\tau)} f(\tau) d\tau. \end{aligned} \quad (14)$$

Let us denote

$$W = \sum_{i=1}^m \alpha_i u(t_i),$$

then (14) implies

$$W = - \sum_{i=1}^m \alpha_i e^{-At_i} W + \sum_{i=1}^m \alpha_i e^{-At_i} u_0 + \sum_{i=1}^m \alpha_i \int_0^{t_i} e^{-A(t_i-\tau)} f(\tau) d\tau,$$

Setting $B = \mathbb{I} + \sum_{i=1}^m \alpha_i e^{-At_i}$, we obtain

$$BW = Bu_0 - u_0 + \sum_{i=1}^m \alpha_i \int_0^{t_i} e^{-A(t_i-\tau)} f(\tau) d\tau,$$

where $\mathbb{I}$ is an identity operator. Condition (10) or (11) guarantee the existence and boundedness of $B^{-1}(A)$, therefore we have

$$W = u_0 - B^{-1}u_0 + B^{-1} \sum_{i=1}^m \alpha_i \int_0^{t_i} e^{-A(t_i-\tau)} f(\tau) d\tau.$$

Now, the equations (13) yield the representation for the solution of (1) in the form $u(t) = u_h(t) + u_{ih}(t)$, where $u_h(t) = e^{-At} B^{-1} u_0$ is the solution of the homogeneous problem with the initial condition u_0 and

$$u_{ih}(t) = -e^{-At} B^{-1} \sum_{i=1}^m \alpha_i \int_0^{t_i} e^{-A(t_i-\tau)} f(\tau) d\tau + \int_0^t e^{-A(t-\tau)} f(\tau) d\tau \quad (15)$$

is the solution of the inhomogeneous problem with zero initial condition.

3 Homogeneous nonlocal problem

First of all we consider the solution $u_h(t) = e^{-At} B^{-1}(A) u_0$ of the homogeneous problem (1). Our aim in this section is to construct an exponentially convergent method for its approximation.

Using the Dunford-Cauchy representation of $u_h(t)$ analogously to [21] we obtain

$$\begin{aligned} u_h(t) &= \frac{1}{2\pi i} \int_{\Gamma_I} e^{-zt} B^{-1}(z) (zI - A)^{-1} u_0 dz \\ &= \frac{1}{2\pi i} \int_{\Gamma_I} \frac{e^{-zt}}{1 + \sum_{i=1}^n \alpha_i e^{-zt_i}} (zI - A)^{-1} u_0 dz. \end{aligned} \quad (16)$$

Representation (16) makes sense only when the function $e^{-zt} B^{-1}(z)$ is analytic in the region enveloped by Γ_I . Let us show, that conditions like (10) or (11) guaranty this analyticity [6].

Actually, the analyticity of $e^{-zt} B^{-1}(z)$ might only be violated when $\sum_{i=1}^n \alpha_i e^{-zt_i} = -1$, since in this case $B^{-1}(z)$ becomes unbounded. It is easy to see that for an arbitrary z we have

$$\begin{aligned} \left| 1 + \sum_{k=1}^m \alpha_k e^{-zt_k} \right| &\geq \left| 1 - \sum_{k=1}^m |\alpha_k| \cdot |e^{t_k a_I \cosh \xi}| \cdot |e^{-it_k b_I \sinh \xi}| \right| \\ &\geq \left| 1 - \sum_{k=1}^m |\alpha_k| e^{-\rho_0 t_k} \right| > 0, \end{aligned}$$

provided that (11) holds true.

Using (8) with $m = 0$ we can modify the representation of $u_h(t)$ as follows:

$$\begin{aligned} u_h(t) &= \frac{1}{2\pi i} \int_{\Gamma_I} e^{-zt} B^{-1}(z) \left[(zI - A)^{-1} - \frac{1}{z} I \right] u_0 dz \\ &= \frac{1}{2\pi i} \int_{\Gamma_I} \frac{e^{-zt}}{1 + \sum_{i=1}^n \alpha_i e^{-zt_i}} \left[(zI - A)^{-1} - \frac{1}{z} I \right] u_0 dz. \end{aligned} \quad (17)$$

After discretization of the integral such modified resolvent provides better convergence speed than (16) in a neighbourhood of $t = 0$ (see [21] for details).

Parameterizing the integral (17) by (5) we get

$$u_h(t) = \frac{1}{2\pi i} \int_{-\infty}^{\infty} \mathcal{F}(t, \xi) d\xi, \quad (18)$$

with

$$\mathcal{F}(t, \xi) = F_A(t, \xi) u_0,$$

$$F_A(t, \xi) = \frac{e^{-z(\xi)t} z'(\xi)}{1 + \sum_{i=1}^n \alpha_i e^{-z(\xi)t_i}} \left[(z(\xi)I - A)^{-1} - \frac{1}{z(\xi)} I \right],$$

$$z'(\xi) = a_I \sinh \xi - ib_I \cosh \xi.$$

Supposing $u_0 \in D(A^\alpha)$, $0 < \alpha < 1$ it was shown in [21] that

$$\begin{aligned} & \|e^{-z(\xi)t} z'(\xi) \left[(z(\xi)I - A)^{-1} - \frac{1}{z(\xi)} I \right] u_0\| \\ & \leq (1 + M) K \frac{b_I}{a_I} \left(\frac{2}{a_I} \right)^\alpha e^{-a_I t \cosh \xi - \alpha |\xi|} \|A^\alpha u_0\|, \quad \xi \in \mathbb{R}, \quad t \geq 0. \end{aligned}$$

The part responsible for the nonlocal condition in (18), can be estimated in the following way

$$\begin{aligned} \left| \left(1 + \sum_{k=1}^m \alpha_k e^{-z(\xi)t_k} \right)^{-1} \right| & \leq \left(1 - \sum_{k=1}^m |\alpha_k| e^{-a_I \cosh(\xi)t_k} \right)^{-1} \\ & \leq \left(1 - \sum_{k=1}^m |\alpha_k| e^{-\rho_1 t_k} \right)^{-1} \equiv Q^{-1}, \end{aligned}$$

where $a_I \geq \rho_1$.

Thus, we obtain the following estimate for $\mathcal{F}(t, \xi)$:

$$\|\mathcal{F}(t, \xi)\| \leq Q(1 + M) K \frac{b_I}{a_I} \left(\frac{2}{a_I} \right)^\alpha e^{-a_I t \cosh \xi - \alpha |\xi|} \|A^\alpha u_0\|, \quad \xi \in \mathbb{R}, \quad t \geq 0. \quad (19)$$

The next step toward a numerical algorithm is an approximation of (18) by the efficient quadrature formula. For this purpose we need to estimate the width of a strip around the real axis where the function $\mathcal{F}(t, \xi)$ permit analytical extension (with respect to ξ). After changing ξ to $\xi + i\nu$ the integration hyperbola Γ_I will be translated into the parametric hyperbola set

$$\begin{aligned} \Gamma(\nu) &= \{z(w) = a_I \cosh(\xi + i\nu) - ib_I \sinh(\xi + i\nu) : \xi \in (-\infty, \infty)\} \\ &= \{z(w) = a(\nu) \cosh \xi - ib(\nu) \sinh \xi : \xi \in (-\infty, \infty)\}, \end{aligned}$$

with

$$\begin{aligned} a(\nu) &= a_I \cos \nu + b_I \sin \nu = \sqrt{a_I^2 + b_I^2} \sin(\nu + \phi/2), \\ b(\nu) &= b_I \cos \nu - a_I \sin \nu = \sqrt{a_I^2 + b_I^2} \cos(\nu + \phi/2), \\ \cos \frac{\phi}{2} &= \frac{b_I}{\sqrt{a_I^2 + b_I^2}}, \quad \sin \frac{\phi}{2} = \frac{a_I}{\sqrt{a_I^2 + b_I^2}}. \end{aligned}$$

The analyticity of the function $\mathcal{F}(t, \xi + i\nu)$, in the strip

$$D_{d_1} = \{(\xi, \nu) : \xi \in (-\infty, \infty), |\nu| < d_1/2\},$$

with some d_1 could be violated if the resolvent or the part related to the nonlocal condition become unbounded. To avoid this we have to choose d_1 in a way that for $\nu \in (-d_1/2, d_1/2)$ the hyperbola set $\Gamma(\nu)$ remains in the right half-plane of the complex plane. For $\nu = -d_1/2$ the corresponding hyperbola is going through the point $(\rho_1, 0)$, for some $0 \leq \rho_1 < \rho_0$. For $\nu = d_1/2$ it coincides with the spectral hyperbola and therefore for all $\nu \in (-d_1/2, d_1/2)$ the set $\Gamma(\nu)$ does not intersect the spectral sector. This fact justifies the choice the hyperbola $\Gamma(0) = \Gamma_I$ as the integration path.

The requirements above imply the following system of equations

$$\begin{cases} a_I \cos(d_1/2) + b_I \sin(d_1/2) = \rho_0, \\ b_I \cos(d_1/2) - a_I \sin(d_1/2) = b_0 = \rho_0 \tan \varphi, \\ a_I \cos(-d_1/2) + b_I \sin(-d_1/2) = \rho_1, \end{cases}$$

it lead us to the next system

$$\begin{cases} a_I = \rho_0 \cos(d_1/2) - b_0 \sin(d_1/2), \\ b_I = \rho_0 \sin(d_1/2) + b_0 \cos(d_1/2), \\ 2a_I \cos(d_1/2) = \rho_0 + \rho_1. \end{cases}$$

Eliminating a_I from the first and the third equations we get

$$\rho_0 \cos d_1 - b_0 \sin d_1 = \rho_1,$$

$$\cos(d_1 + \varphi) = \frac{\rho_1}{\sqrt{\rho_0^2 + b_0^2}},$$

i.e.

$$d_1 = \arccos\left(\frac{\rho_1}{\sqrt{\rho_0^2 + b_0^2}}\right) - \varphi, \quad (20)$$

with $\cos \varphi = \frac{\rho_0}{\sqrt{\rho_0^2 + b_0^2}}$, $\sin \varphi = \frac{b_0}{\sqrt{\rho_0^2 + b_0^2}}$. Thus, for a_I, b_I we receive

$$\begin{aligned}
a_I &= \sqrt{\rho_0^2 + b_0^2} \cos \left(\frac{d_1}{2} + \varphi \right) \\
&= \rho_0 \frac{\cos \left(\frac{d_1}{2} + \varphi \right)}{\cos \varphi} = \rho_0 \frac{\cos \left(\arccos \left(\frac{\rho_1}{\sqrt{\rho_0^2 + b_0^2}} \right) / 2 + \varphi / 2 \right)}{\cos \varphi}, \\
b_I &= \sqrt{\rho_0^2 + b_0^2} \sin \left(\frac{d_1}{2} + \varphi \right) \\
&= \rho_0 \frac{\sin \left(\frac{d_1}{2} + \varphi \right)}{\cos \varphi} = \rho_0 \frac{\sin \left(\arccos \left(\frac{\rho_1}{\sqrt{\rho_0^2 + b_0^2}} \right) / 2 + \varphi / 2 \right)}{\cos \varphi}.
\end{aligned} \tag{21}$$

For a_I and b_I defined as above the vector valued function $\mathcal{F}(t, w)$ is analytic in the strip D_{d_1} with respect to $w = \xi + i\nu$ for any $t \geq 0$. Note, that for $\rho_1 = 0$ we have $d_1 = \pi/2 - \varphi$ as in [21].

Taking into account (21) we similarly can write equations for $a(\nu), b(\nu)$ on the whole interval $-\frac{d_1}{2} \leq \nu \leq \frac{d_1}{2}$

$$\begin{aligned}
a(\nu) &= a_I \cos \nu + b_I \sin \nu = \sqrt{\rho_0^2 + b_0^2} \cos \left(\frac{d_1}{2} + \varphi \right) \cos(\nu) \\
&\quad + \sqrt{\rho_0^2 + b_0^2} \sin \left(\frac{d_1}{2} + \varphi \right) \sin(\nu) = \sqrt{\rho_0^2 + b_0^2} \cos \left(\frac{d_1}{2} + \varphi - \nu \right), \\
b(\nu) &= b_I \cos \nu - a_I \sin \nu = \sqrt{\rho_0^2 + b_0^2} \sin \left(\frac{d_1}{2} + \varphi \right) \cos(\nu) \\
&\quad - \sqrt{\rho_0^2 + b_0^2} \cos \left(\frac{d_1}{2} + \varphi \right) \sin(\nu) = \sqrt{\rho_0^2 + b_0^2} \sin \left(\frac{d_1}{2} + \varphi - \nu \right), \\
\rho_1 &\leq a(\nu) \leq \rho_0, \quad b_0 \leq b(\nu) \leq \sqrt{b_0^2 + \rho_0^2 - \rho_1^2},
\end{aligned}$$

with d_1 , defined by (20).

For the part responsible for the nonlocal condition we have

$$\begin{aligned}
\left| \left(1 + \sum_{i=1}^n \alpha_i e^{-z(\xi, \nu) t_i} \right)^{-1} \right| &\leq \left(1 - \sum_{i=1}^n |\alpha_i| |e^{-z(\xi, \nu) t_i}| \right)^{-1} \\
&\leq \left(1 - \sum_{i=1}^n |\alpha_i| e^{-a(\nu) \cosh(\xi) t_i} \right)^{-1}
\end{aligned}$$

$$\leq \left(1 - \sum_{i=1}^n |\alpha_i| e^{-a(\nu)t_i}\right)^{-1} < \left(1 - \sum_{i=1}^n |\alpha_i| e^{-\rho_1 t_i}\right)^{-1} < q^{-1} = Q,$$

where $z(\xi, \nu) = a(\nu) \cosh(\xi) - ib(\nu) \sinh(\xi)$.

Now, for $w \in D_{d_1}$ we get the estimate

$$\begin{aligned} \|\mathcal{F}(t, w)\| &\leq e^{-a(\nu)t \cosh \xi} \frac{(1+M)QK \sqrt{a^2(\nu) \sinh^2 \xi + b^2(\nu) \cosh^2 \xi}}{(a^2(\nu) \cosh^2 \xi + b^2(\nu) \sinh^2 \xi)^{(1+\alpha)/2}} \|A^\alpha u_0\| \\ &\leq (1+M)QK \frac{b(\nu)}{a(\nu)} \frac{e^{-a(\nu)t \cosh \xi}}{(a^2(\nu) \cosh^2 \xi + b^2(\nu) \sinh^2 \xi)^{(\alpha/2)}} \|A^\alpha u_0\| \\ &\leq (1+M)QK \frac{b(\nu)}{a(\nu)} \left(\frac{2}{a(\nu)}\right)^\alpha e^{-a(\nu)t \cosh \xi - \alpha|\xi|} \|A^\alpha u_0\| \\ &\leq (1+M)QK \tan\left(\frac{d_1}{2} + \varphi - \nu\right) \left(\frac{2 \cos \varphi}{\rho_0 \cos\left(\frac{d_1}{2} + \varphi - \nu\right)}\right)^\alpha e^{-\alpha|\xi|} \|A^\alpha u_0\|, \\ &\quad \forall w \in D_d. \end{aligned}$$

Similarly to [23], we introduce the space $\mathbf{H}^p(D_d)$, $1 \leq p \leq \infty$ of all vector-valued functions $\mathcal{F}$ analytic in the strip

$$D_d = \{z \in \mathbb{C} : -\infty < \Re z < \infty, |\Im z| < d\},$$

equipped by the norm

$$\|\mathcal{F}\|_{\mathbf{H}^p(D_d)} = \begin{cases} \lim_{\epsilon \rightarrow 0} (\int_{\partial D_d(\epsilon)} \|\mathcal{F}(z)\|^p |dz|)^{1/p} & \text{if } 1 \leq p < \infty, \\ \lim_{\epsilon \rightarrow 0} \sup_{z \in \partial D_d(\epsilon)} \|\mathcal{F}(z)\| & \text{if } p = \infty, \end{cases}$$

where

$$D_d(\epsilon) = \{z \in \mathbb{C} : |\Re(z)| < 1/\epsilon, |\Im(z)| < d(1 - \epsilon)\}$$

and $\partial D_d(\epsilon)$ is the boundary of $D_d(\epsilon)$.

Taking into account that the integrals over the vertical sides of the rectangle $D_{d_1}(\epsilon)$ vanish as $\epsilon \rightarrow 0$, the above estimate for $\|\mathcal{F}(t, w)\|$ implies

$$\begin{aligned} \|\mathcal{F}(t, \cdot)\|_{\mathbf{H}^1(D_{d_1})} &\leq \|A^\alpha u_0\| [C_-(\varphi, \alpha) \\ &\quad + C_+(\varphi, \alpha)] \int_{-\infty}^{\infty} e^{-\alpha|\xi|} d\xi = C(\varphi, \alpha) \|A^\alpha u_0\| \end{aligned} \tag{22}$$

with

$$\begin{aligned} C(\varphi, \alpha) &= \frac{2}{\alpha} [C_+(\varphi, \alpha) + C_-(\varphi, \alpha)], \\ C_\pm(\varphi, \alpha) &= (1+M)QK \tan\left(\frac{d_1}{2} + \varphi \pm \frac{d_1}{2}\right) \left(\frac{2 \cos \varphi}{\rho_0 \cos\left(\frac{d_1}{2} + \varphi \pm \frac{d_1}{2}\right)}\right)^\alpha. \end{aligned}$$

Note that the influence of both the smoothness parameter of u_0 given by α and of the spectral characteristics of the operator A given by φ and ρ_0 is accounted by that fact, that the constant $C(\varphi, \alpha)$ from (21) tends to ∞ if $\alpha \rightarrow 0$, $\varphi \rightarrow \pi/2$ or $\rho_1 \rightarrow 0$ (in this case due to (20) $d_1 \rightarrow \frac{\pi}{2} - \varphi$).

We approximate integral (18) by the following Sinc-quadrature [23, 21]:

$$u_{h,N}(t) = \frac{h}{2\pi i} \sum_{k=-N}^N \mathcal{F}(t, z(kh)), \quad (23)$$

with an error

$$\begin{aligned} \|\eta_N(\mathcal{F}, h)\| &= \|u_h(t) - u_{h,N}(t)\| \\ &\leq \|u_h(t) - \frac{h}{2\pi i} \sum_{k=-\infty}^{\infty} \mathcal{F}(t, z(kh))\| + \|\frac{h}{2\pi i} \sum_{|k|>N} \mathcal{F}(t, z(kh))\| \\ &\leq \frac{1}{2\pi} \frac{e^{-\pi d_1/h}}{2 \sinh(\pi d_1/h)} \|\mathcal{F}\|_{\mathbf{H}^1(D_{d_1})} \\ &\quad + \frac{C(\varphi, \alpha)h \|A^\alpha u_0\|}{2\pi} \sum_{k=N+1}^{\infty} \exp[-a_I t \cosh(kh) - \alpha kh] \\ &\leq \frac{c \|A^\alpha u_0\|}{\alpha} \left\{ \frac{e^{-\pi d_1/h}}{\sinh(\pi d_1/h)} + \exp[-a_I t \cosh((N+1)h) - \alpha(N+1)h] \right\}, \end{aligned}$$

where the constant c does not depend on h, N, t . Equalizing the both exponentials for $t = 0$ implies

$$\frac{2\pi d_1}{h} = \alpha(N+1)h,$$

or after the transformation

$$h = \sqrt{\frac{2\pi d_1}{\alpha(N+1)}}. \quad (24)$$

With this step-size the following error estimate holds true

$$\|\eta_N(\mathcal{F}, h)\| \leq \frac{c}{\alpha} \exp\left(-\sqrt{\frac{\pi d_1 \alpha}{2}}(N+1)\right) \|A^\alpha u_0\|, \quad (25)$$

with a constant c independent of t, N . In the case $t > 0$ the first summand in the argument of $\exp[-a_I t \cosh((N+1)h) - \alpha(N+1)h]$ from the estimate for $\|\eta_N(\mathcal{F}, h)\|$ contributes mainly to the error order. Setting in this case

$h = c_1 \ln N/N$ with some positive constant c_1 we remain, asymptotically for a fixed t , with an error

$$\|\eta_N(\mathcal{F}, h)\| \leq c [e^{-\pi d_1 N/(c_1 \ln N)} + e^{-c_1 a_I t N/2 - c_1 \alpha \ln N}] \|A^\alpha u_0\|, \quad (26)$$

where c is a positive constant. Thus, we have proved the following result.

Theorem 2 *Let A be a densely defined strongly positive operator and $u_0 \in D(A^\alpha)$, $\alpha \in (0, 1)$, then the Sinc-quadrature (23) represents an approximate solution of the homogeneous nonlocal value problem (1) (i.e. the case when $f(t) \equiv 0$) and possesses an exponential convergence rate which is uniform with respect to $t \geq 0$ and is of the order $\mathcal{O}(e^{-c\sqrt{N}})$ uniformly in $t \geq 0$ provided that $h = \mathcal{O}(1/\sqrt{N})$ (estimate (25)) and of the order $\mathcal{O}(\max\{e^{-\pi d N/(c_1 \ln N)}, e^{-c_1 a_I t N/2 - c_1 \alpha \ln N}\})$ for each fixed $t > 0$ provided that $h = c_1 \ln N/N$ (estimate (26)).*

4 Inhomogeneous nonlocal problem

In this section we consider the particular solution (15) of inhomogeneous problem (1), i.e. with $f(t) \neq 0$.

Let us rewrite formula (15) in the form

$$u_{ih}(t) = u_{1,ih}(t) + u_{2,ih}(t), \quad (27)$$

with

$$u_{1,ih}(t) = \int_0^t e^{-A(t-\tau)} f(\tau) d\tau, \quad u_{2,ih}(t) = - \sum_{j=1}^m \alpha_j u_{2,ih,j}(t), \quad (28)$$

where

$$u_{2,ih,j}(t) = \int_0^{t_j} B^{-1} e^{-A(t+t_j-\tau)} f(\tau) d\tau. \quad (29)$$

We approximate the term $u_{1,ih}(t)$ by the algorithm proposed in [21]:

$$\begin{aligned} u_{1,ih}(t) \approx u_{1,N}(t) &= \frac{h}{2\pi i} \sum_{k=-N}^N z'(kh) [(z(kh)I - A)^{-1} - \frac{1}{z(kh)} I] \\ &\times h \sum_{p=-N}^N \mu_{k,p}(t) f(\omega_p(t)), \end{aligned} \quad (30)$$

where

$$\mu_{k,p}(t) = \frac{t}{2} \exp\left\{-\frac{t}{2} z(kh) [1 - \tanh(ph)]\right\} / \cosh^2(ph),$$

$$\omega_p(t) = \frac{t}{2}[1 + \tanh(ph)], \quad h = \mathcal{O}\left(1/\sqrt{N}\right),$$

$$z(\xi) = a_I \cosh \xi - ib_I \sinh \xi, \quad z'(\xi) = a_I \sinh \xi - ib_I \cosh \xi.$$

The next theorem (see [21]) characterizes the error of this algorithm.

Theorem 3 *Let A be a densely defined strongly positive operator with spectral characteristics ρ_0 , φ and a right hand side $f(t) \in D(A^\alpha)$, $\alpha > 0$ for $t \in [0, \infty]$ can be analytically extended into the sector $\Sigma_f = \{\rho e^{i\theta_1} : \rho \in [0, \infty], |\theta_1| < \varphi\}$ where the estimate*

$$\|A^\alpha f(w)\| \leq c_\alpha e^{-\delta_\alpha |Re w|}, \quad w \in \Sigma_f \quad (31)$$

with $\delta_\alpha \in (0, \sqrt{2}\rho_0]$ holds, then algorithm (30) converges to the solution of (1) with the error estimate

$$\|\mathcal{E}_N(t)\| = \|u_{1,ih}(t) - u_{1,N}(t)\| \leq ce^{-c_1\sqrt{N}} \quad (32)$$

uniformly in t with positive constants c , c_1 depending on α , φ , ρ_0 and independent of N .

Let us construct an exponentially convergent approximation to the term $u_{2,ih}$. Using the representation of the operator functions by means of the Dunford-Cauchy integral for the j -th summand of $u_{2,ih}$ we get

$$\begin{aligned} u_{2,ih,j}(t) &= \int_0^{t_j} \frac{1}{2\pi i} \int_{\Gamma_I} e^{-z(t+t_j-s)} B^{-1}(z) [(zI - A)^{-1} - \frac{1}{z}I] f(s) dz ds \\ &= \frac{1}{2\pi i} \int_{\Gamma_I} e^{-z(\xi)t} B^{-1}(z) \left[(z(\xi)I - A)^{-1} - \frac{1}{z(\xi)}I \right] \\ &\quad \times \int_0^{t_j} e^{-z(\xi)(t_j-s)} f(s) ds z'(\xi) d\xi, \\ z(\xi) &= a_I \cosh \xi - ib_I \sinh \xi. \end{aligned} \quad (33)$$

Replacing here the first integral by the Sinc-quadrature with $h = \mathcal{O}(N^{-1/2})$ we obtain

$$\begin{aligned} u_{2,ih,j}(t) \approx u_{2,N}(t) &= \frac{h}{2\pi i} \sum_{k=-N}^N e^{-z(kh)t} z'(kh) B^{-1}(z(kh)) \\ &\quad \times \left[(z(kh)I - A)^{-1} - \frac{1}{z(kh)}I \right] f_{k,j}, \end{aligned} \quad (34)$$

where

$$f_{k,j} = \int_0^{t_j} e^{-z(kh)(t_j-s)} f(s) ds, \quad k = -N, \dots, N, \quad j = \overline{1, n}. \quad (35)$$

In order to construct an exponentially convergent quadrature for these integrals we change the variables by

$$s = \frac{t_j}{2}(1 + \tanh \xi), \quad (36)$$

and obtain the improper integrals

$$f_{k,j} = \int_{-\infty}^{\infty} \mathcal{F}_k(t_j, \xi) d\xi, \quad (37)$$

with

$$\mathcal{F}_k(t_j, \xi) = \frac{t_j}{2 \cosh^2 \xi} \exp[-z(kh)t_j(1 - \tanh \xi)/2] f(t_j(1 + \tanh \xi)/2). \quad (38)$$

Note that equation (36) represents the conformal mapping $w = \psi(z) = t_j[1 + \tanh z]/2$, $z = \phi(w) = \frac{1}{2} \ln \frac{t_j - w}{w}$ of the strip D_ν onto the eye-shaped domain $\mathcal{A}_\nu$ [23]. On the real axes the integrand can be estimated by

$$\begin{aligned} \|\mathcal{F}_k(t_j, \xi)\| &\leq \frac{t_j}{2 \cosh^2 \xi} \exp[-a_I \cosh(kh)t_j(1 - \tanh \xi)/2] \\ &\quad \times \|f(t_j(1 + \tanh \xi)/2)\| \\ &\leq 2t_j e^{-2|\xi|} \|f(t_j(1 + \tanh \xi)/2)\|. \end{aligned} \quad (39)$$

The following result from [21] is needed to justify a Sinc-quadrature for (37):

Lemma 4 *Let the right-hand side $f(t)$ in (1) for $t \in [0, \infty]$ can be analytically extended into the sector $\Sigma_f = \{\rho e^{i\theta_1} : \rho \in [0, \infty], |\theta_1| < \varphi\}$ and for all complex $w \in \Sigma_f$ we have*

$$\|f(w)\| \leq c e^{-\delta |\operatorname{Re} w|}, \quad (40)$$

with $\delta \in (0, \sqrt{2}\rho_0]$, then the integrand $\mathcal{F}_k(t_j, \xi)$ can be analytically extended into a strip D_{d_1} , $0 < d_1 < \varphi/2$ and belongs to the class $\mathbf{H}^1(D_{d_1})$ with respect to ξ , where ρ_0, φ are the spectral characteristics of A .

Under the assumptions of Lemma 4 we can construct the following exponentially convergent Sinc-approximation to $f_{k,j}$:

$$f_{k,j} \approx f_{k,j,N} = h \sum_{p=-N}^N \mu_{k,p,j} f(\omega_{p,j}), \quad (41)$$

where

$$\begin{aligned}\mu_{k,p,j} &= \frac{t_j}{2} \frac{\exp\{-\frac{t_j}{2}z(kh)[1 - \tanh(ph)]\}}{\cosh^2(ph)}, \\ \omega_{p,j} &= \frac{t_j}{2}[1 + \tanh(ph)], \quad h = \mathcal{O}(1/\sqrt{N}), \\ z(\xi) &= a_I \cosh \xi - ib_I \sinh \xi.\end{aligned}$$

Substituting (41) into (34) we get the following algorithm to compute an approximation $u_{2,j,N}$ to $u_{2,ih,j}$

$$\begin{aligned}u_{2,ih,j}(t) \approx u_{2,j,N}(t) &= \frac{h}{2\pi i} \sum_{k=-N}^N e^{-z(kh)t} z'(kh) B^{-1}(z(kh)) \\ &\times \left[(z(kh)I - A)^{-1} - \frac{1}{z(kh)} I \right] h \sum_{p=-N}^N \mu_{k,p,j} f(\omega_{p,j}).\end{aligned}\tag{42}$$

We represent the error in the form

$$\mathcal{E}_N(t) = u_{2,ih,j}(t) - u_{2,j,N}(t) = r_{1,N}(t) + r_{2,N}(t),\tag{43}$$

where

$$\begin{aligned}r_{1,N}(t) &= u_{2,ih,j}(t) - u_{2,N}(t), \\ r_{2,N}(t) &= u_{2,N}(t) - u_{2,j,N}(t).\end{aligned}\tag{44}$$

Using estimate (25)(see also Theorem 2) we obtain the estimate for $r_{1,N}(t)$

$$\begin{aligned}\|r_{1,N}(t)\| &= \left\| \int_0^{t_j} \left\{ \frac{1}{2\pi i} \int_{-\infty}^{\infty} F_A(t + t_j - s, \xi) d\xi \right. \right. \\ &\quad \left. \left. - \frac{h}{2\pi i} \sum_{k=-N}^N F_A(t + t_j - s, kh) \right\} f(s) ds \right\| \\ &\leq \frac{c}{\alpha} \exp\left(-\sqrt{\frac{\pi d\alpha}{2}}(N+1)\right) \int_0^{t_j} \|A^\alpha f(s)\| ds,\end{aligned}\tag{45}$$

where $F_A(t, \xi)$ is the operator defined in section 3. Due to (9) for $m = 0$ we have for the error $r_{2,N}$

$$\begin{aligned}\|r_{2,N}(t)\| &= \left\| \frac{h}{2\pi i} \sum_{k=-N}^N e^{-z(kh)t} z'(kh) B(z(kh)) \right. \\ &\quad \left. \times \left[(z(kh)I - A)^{-1} - \frac{1}{z(kh)} I \right] R_{k,j} \right\| \\ &\leq \frac{h(1+M)QK}{2\pi} \sum_{k=-N}^N \frac{|e^{-z(kh)t} z'(kh)|}{|z(kh)|^{1+\alpha}} \|A^\alpha R_{k,j}\|,\end{aligned}\tag{46}$$

where

$$R_{k,j} = f_{k,j} - f_{k,j,N}.$$

The estimate (39) yields

$$\|A^\alpha \mathcal{F}_k(t_j, \xi)\| \leq 2t_j e^{-2|\xi|} \|A^\alpha f(\frac{t_j}{2}(1 + \tanh \xi))\|. \quad (47)$$

Due to Lemma 4 the assumption $\|A^\alpha f(w)\| \leq c_\alpha e^{-\delta_\alpha |\operatorname{Re} w|} \forall w \in \Sigma_f$ guarantees that $A^\alpha f(w) \in \mathbf{H}^1(D_{d_1})$ and $A^\alpha \mathcal{F}_k(t_j, w) \in \mathbf{H}^1(D_{d_1})$. These two conditions turn us to the situation as in proposition of Theorem 3.2.1, p.144 from [23] with $A^\alpha f(w)$ instead of f which implies

$$\begin{aligned} \|A^\alpha R_{k,j}\| &= \|A^\alpha (f_{k,j} - f_{k,j,N})\| \\ &= \left\| \int_{-\infty}^{\infty} A^\alpha \mathcal{F}_k(t_j, \xi) d\xi - h \sum_{k=-\infty}^{\infty} A^\alpha \mathcal{F}_k(t_j, kh) \right\| + \left\| h \sum_{|k|>N} A^\alpha \mathcal{F}_k(t_j, kh) \right\| \\ &\leq \frac{e^{-\pi d_1/h}}{2 \sinh(\pi d_1/h)} \|\mathcal{F}_k(t_j, w)\|_{\mathbf{H}^1(D_{d_1})} \\ &\quad + h \sum_{|k|>N} 2t_j e^{-2|kh|} \|A^\alpha f(\frac{t_j}{2}(1 + \tanh kh))\| \\ &\leq c e^{-2\pi d_1/h} \|A^\alpha f(t_j, w)\|_{\mathbf{H}^1(D_{d_1})} \\ &\quad + h \sum_{|k|>N} 2t_j e^{-2|kh|} c_\alpha \exp \left\{ -\delta_\alpha \frac{t_j}{2}(1 + \tanh kh) \right\}, \end{aligned}$$

i.e., we have

$$\|A^\alpha R_{k,j}\| \leq c e^{-c_1 \sqrt{N}}, \quad (48)$$

where positive constants $c_\alpha, \delta_\alpha, c, c_1$ do not depend on t, N, k . Now, (46) takes the form

$$\begin{aligned} \|r_{2,N}(t)\| &= \frac{h}{2\pi i} \sum_{k=-N}^N e^{-z(kh)t} z'(kh) B(z(kh)) \\ &\quad \times \left[(z(kh)I - A)^{-1} - \frac{1}{z(kh)} I \right] R_{k,j} \\ &\leq c e^{-c_1 \sqrt{N}} S_N(t), \end{aligned} \quad (49)$$

with $S_N(t) = \sum_{k=-N}^N h \frac{|e^{-z(kh)t} z'(kh)|}{|z(kh)|^{1+\alpha}}$. Using the estimate (4.8) from [21] and

$$\begin{aligned} |z(kh)| &= \sqrt{a_I^2 \cosh^2(kh) + b_I^2 \sinh^2(kh)} \\ &\geq a_I \cosh(kh) \geq a_I e^{|kh|}/2, \end{aligned} \quad (50)$$

the last sum can be estimated by

$$|S_N(t)| \leq \frac{c}{\sqrt{N}} \sum_{k=-N}^N e^{-\alpha|k|/\sqrt{N}} \leq c \int_{-\sqrt{N}}^{\sqrt{N}} e^{-\alpha t} dt \leq c/\alpha \quad \forall t \in [0, \infty). \quad (51)$$

Taking into account (48), (51) we deduce from (49)

$$\|r_{2,N}(t)\| \leq ce^{-c_1\sqrt{N}}. \quad (52)$$

The next assertion now follows from (43), (45) and (52).

Theorem 5 *Let the assumptions of theorem 3 hold. Then algorithm (42) converges uniformly with respect to t and moreover the following error estimate holds true:*

$$\|\mathcal{E}_N(t)\| = \|u_{2,ih,j}(t) - u_{2,j,N}(t)\| \leq ce^{-c_1\sqrt{N}}, \quad (53)$$

with positive constants c, c_1 depend on α, φ, ρ_0 and independent of N .

Now we can use approximation (30) for the every summand and get:

$$u_{2,ih}(t) = \sum_{i=1}^n \alpha_i \int_0^{t_i} B^{-1} e^{-A(t+t_i-\tau)} f(\tau) d\tau \approx \sum_{j=1}^n u_{2,j,N}(t) = u_{2,N}(t). \quad (54)$$

Theorem 5 guarantees that the error this approach will be bounded by:

$$\|u_{2,ih}(t) - u_{2,N}(t)\| \leq \sum_{j=1}^m ce^{-c_1\sqrt{N}} \leq c_2 e^{-c_1\sqrt{N}}. \quad (55)$$

Thus, the approximations (23) together with (30) and (54) represent an exponentially convergent algorithm for the problem (1).

5 Numerical examples

Example 6 *We consider the homogeneous problem (1) with the operator A defined by*

$$\begin{aligned} D(A) &= \{u(x) \in H^2(0, 1) : u(0) = u(1) = 0\}, \\ Au &= -u''(x) \quad \forall u \in D(A). \end{aligned} \quad (56)$$

The initial nonlocal condition reads as follows:

$$u(x, 0) + 0.5u(x, 0.2) + 0.3u(x, 0.4) = (1 + 0.5e^{-\pi^2 0.2} + 0.3e^{-\pi^2 0.4}) \sin(\pi x),$$

with $u_0 = (1 + 0.5e^{-\pi^2 0.2} + 0.3e^{-\pi^2 0.4}) \sin(\pi x) \in D(A)$. The exact solution of the problem is $u(x, t) = e^{-\pi^2 t} \sin(\pi x)$. It is easy to find that

$$(zI - A)^{-1}u_0 = \left(z + \frac{d^2}{dx^2}\right)^{-1} \sin(\pi x) = \frac{\sin(\pi x)}{z - \pi^2}.$$

Calculations with the algorithm (42) above has been performed in Maple with $h = N^{-1/2}$. The error at $x = 0.5$, $t = 0.3$ is presented in Table 1 and clearly exhibits an exponential decay according to the theoretical estimates.

N	ε_N
4	.29857983847712589e-1
8	.41823888073604986e-2
16	.11258594468208641e-2
32	.10042178166563831e-3
64	.28007158539828452e-5
128	.2098826601399176e-7
256	.1858929920173152e-10
512	.856837124351510e-15

Table 1: The error for $x = 0.5$, $t = 0.3$.

Due to Theorem 2 the error should not be greater then $\varepsilon_N = \mathcal{O}\left(e^{-c\sqrt{N}}\right)$. The constant c in the exponent can be estimated using the following a-posteriori relation:

$$c = \ln\left(\frac{\varepsilon_N}{\varepsilon_{2N}}\right) (\sqrt{2} - 1)^{-1} N^{-1/2} = \ln(\mu_N) (\sqrt{2} - 1)^{-1} N^{-1/2}.$$

The numerical results are presented in the Table 2 for this estimation show that the constant can be estimated as $c \approx 1.5$ when $N \rightarrow \infty$.

The next example deals again with a homogeneous problem but in this more realistic case the resolvent of A on the element u_0 cannot be calculated analytically.

Example 7 We consider the homogeneous problem (1) with the operator A defined as in (56) and with the following initial nonlocal condition:

$$u(x, 0) + u(x, 0.5) = x \ln(x),$$

N	c
4	2.372652515388745588587496
8	1.120148732795449515627946
16	1.458741976765153165445005
32	1.527648924601130131250452
64	1.476794596387591759032900
128	1.499935011373075736075927
256	1.506597339081609844717370

Table 2: The estimate of c

where $u_0 = x \ln(x) \in A^\alpha$, $\alpha < 1/2$. In this case the resolvent can be represented using the Green function

$$(zI - A)^{-1}u_0 = \left(z + \frac{d^2}{dx^2}\right)^{-1} x \ln(x) = \int_0^1 G(x, s) s \ln(s) ds,$$

$$G(x, s) = -\frac{1}{\sqrt{z} \sin(\sqrt{z})} \begin{cases} \sin(x\sqrt{z}) \sin((1-s)\sqrt{z}) & x \leq s, \\ \sin(s\sqrt{z}) \sin((1-x)\sqrt{z}) & x \geq s \end{cases},$$

where the integrals were computed by exponentially convergent Sinc-quadrature (see e.g. [23]) using Maple. The results for $x = 0.5$, $t = 0.3$ are presented in Table 3.

N	$u(x, t)$
4	-.241535790017043e-1
8	-.228401191029108e-1
16	-.194273285627507e-1
32	-.192905848633180e-1
64	-.192911920318628e-1
128	-.192907849909929e-1
256	-.192907820740651e-1

Table 3: Values of the solution $u(x, t)$ for $x = 0.5$, $t = 0.3$.

It can be easily seen that the number of stabilized digits increases according to the theoretical prediction by Theorem 2.

Example 8 Let us consider the inhomogeneous problem (1) with the same A defined by (56), and the nonlocal condition

$$u(x, 0) + 0.5u(x, 0.2) = (1 + 0.5e^{0.2}) \sin(\pi x),$$

For $f(t, x)$ at the right-hand side of the equation (1) we set

$$f(x, t) = (1 + \pi^2)e^t \sin(\pi x).$$

The exact solution of the problem is $u(x, t) = e^t \sin(\pi x)$. We have used the algorithm defined by (23), (30), (54) and implemented in Maple. For $x = 0.5$, $t = 0.3$ the results presented in Table 4 and are again in good agreement with the theoretical predictions.

N	ε_N
4	.202211483120243
8	.726677678737409e-1
16	.138993889900620e-1
32	.143037059411419e-2
64	.554542099757830e-4
128	.532640823981411e-6
256	.730569324317506e-9
512	.648376079810788e-13

Table 4: The error for $x = 0.5$, $t = 0.3$.

Acknowledgment. The authors would like to acknowledge the support provided by the Deutsche Forschungsgemeinschaft (DFG).

References

- [1] A.A. Samarskii (1980). Some problems of the theory of differential equations. *Differentsial'nye Uravneniya* (Differential equations). 16, No. 11:1925–1935.
- [2] A.W. Leung, and G.-S. Chen (1999). Optimal Control of Multigroup Neutron Fission Systems. *J. Appl. Math. and Opt.* 40, No. 1:39–60.
- [3] A.W. Leung, and L.A. Ortega (1998). Existence and monotone scheme for time-periodic nonquasimonotone reaction-diffusion systems: application to autocatalytic chemistry. *J. Math. Anal. Appl.* 221, No. 2:712–733.
- [4] D. Gordeziani, and G. Avalishvili (2001). Investigation of the nonlocal initial boundary value problems for some hyperbolic equations. *Hiroshima Math. J.* 31, No.3:345–366.

- [5] Ph. Clement, H.J.A.M. Heijmans, S. Angenent, C.J. van Duijn, and B. de Pagter (1987). One-parameter semigroups. CWI Monographs, 5. North-Holland Publishing Co. Amsterdam.
- [6] S.G. Krein (1971). Linear Differential Operators in Banach Spaces. Amer. Math. Soc. New York.
- [7] L. Byszewski (1991). Theorems about the existence and uniqueness of solution of a semilinear evolution nonlocal Cauchy problem. Journal of Mathematical analysis and Applications. 162, No.5:494–505.
- [8] L. Byszewski (1998). Application of properties of the right-hand sides of evolution equations to an investigation of nonlocal evolution problems. Nonlinear Anal. 33, No.5:413–426.
- [9] M.L. Fernandez, Ch. Lubich, C. Palencia, and A. Schädle (2005). Fast Runge-Kutta approximation of inhomogeneous parabolic equations. Numerische Mathematik. 5:1-17.
- [10] I.P. Gavriljuk, W. Hackbusch, and B.N. Khoromskij (2005). Data-sparse approximation of a class of operator-valued functions. Math. Comp. 74:681–708.
- [11] I.P. Gavriljuk, W. Hackbusch, and B.N. Khoromskij (2005). Tensor-Product Approximation to Elliptic and Parabolic Solution Operators in Higher Dimensions. Computing. 74:131–157.
- [12] I.P. Gavriljuk, and V.L. Makarov (2001). Exponentially convergent parallel discretization methods for the first order evolution equations. Computational Methods in Applied Mathematics (CMAM) 1, No.4:333- 355.
- [13] M. López-Fernández, and C. Palencia (2004). On the numerical inversion of the Laplace transform of certain holomorphic mappings. Applied Numerical Mathematics. 51:289–303.
- [14] M. López-Fernández, C. Palencia, and A. Schädle (2006). A spectral order method for inverting sectorial Laplace transforms. SIAM J. Numer. Anal. 44:1332–1350.
- [15] D. Sheen, I.H. Sloan, and V. Thomée (2000). A parallel method for time-discretization of parabolic problems based on contour integral representation and quadrature. Math.Comp.69:177–195.

- [16] D. Sheen, I.H. Sloan, and V. Thomée (2003). A parallel method for time-discretization of parabolic equations based on Laplace transformation and quadrature. *IMA Journal of Numerical Analysis*. 23:269–299.
- [17] V. Vasylyk (2003). Uniform exponentially convergent method for the first order evolution equation with unbounded operator coefficient. *Journal of Numerical and Applied Mathematics* (ISSN 0868-6912). 1:99–104 (in Russian).
- [18] V. Vasylyk (2004). Approximate solution of the Cauchy problem for differential equation with sectorial operator in Banach space. *Bulletin of Lviv University. Physical and Mathematical sciences*. 9:34–46 (in Ukrainian).
- [19] J. A. C. Weideman (2006). Optimizing Talbot’s contours for the inversion of the Laplace transform. *SIAM J. Numer. Anal.* 44, No.6:2342–2362.
- [20] J. A. C. Weideman, and L. N. Trefethen (2007). Parabolic and hyperbolic contours for computing the Bromwich integral. *Math. Comp.* 76, No.256:1341–1356.
- [21] I.P. Gavrilyuk, and V.L. Makarov (2005). Exponentially convergent algorithms for the operator exponential with applications to inhomogeneous problems in Banach spaces. *SIAM Journal on Numerical Analysis*. 43, No.5:2144–2171.
- [22] K. Yosida (1968). *Functional Analysis*. Springer-Verlag.
- [23] F. Stenger (1993). *Numerical methods based on Sinc and analytic functions*. Springer Verlag. New York, Berlin, Heidelberg.